

Some examples in the integral and Brown-Peterson cohomology of p -groups.

I. J. Leary

School of Mathematical Sciences,
Queen Mary and Westfield College,
London.
E1 4NS

N. Yagita

Musashi Institute of Technology,
Tamazatumi,
Setagaya,
Tokyo.

Introduction.

For a finite group G , we define the Chern ring, $\text{Ch}(G)$, to be the subring of $H^{\text{even}}(G; \mathbb{Z})$ generated by Chern classes of representations of G . We say that G has p -rank n if n is maximal such that G contains a subgroup isomorphic to $(C_p)^n$. In [3] Atiyah showed that for any finite group G , $K^0(BG)$ is the completion of the representation ring of G with respect to a certain topology. The filtration of $K^0(BG)$ given by the E_∞ page of the Atiyah-Hirzebruch spectral sequence (AHSS) gives rise to a filtration of the representation ring of G . Atiyah conjectured that this filtration coincided with another filtration defined algebraically, and remarked that this conjecture is equivalent to the conjecture that $\text{Ch}(G)$ maps onto the E_∞ page of the AHSS. (It is clear that $\text{Ch}(G)$ consists of universal cycles because the AHSS for $BU(n)$ collapses.) Weiss discovered that the alternating group A_4 gives a counterexample to this conjecture [16], and Thomas has exhibited many counterexamples, all of which have order divisible by more than one prime [14].

Thomas showed that the split metacyclic p -groups and various other p -groups of p -rank two have the property that the Chern subring is the whole of the even degree integral cohomology, and conjectured that this property would hold for all p -groups of p -rank two [12], [13], [15]. The group A_4 shows that the conjecture cannot be extended to groups of non-prime power order. AlZubaidy claimed to have verified this conjecture, but some of his proofs are flawed [1], [2]. Recently Huebschmann and Tezuka-Yagita have shown that

$\text{Ch}(G) = H^{\text{even}}(G; \mathbb{Z})$ for any metacyclic p -group G [6], [11]. For $p \geq 5$ Blackburn's classification [4] implies that the only p -groups of p -rank two not covered by the above theorems are the groups $G(n, \epsilon)$ defined below. We show that $\text{Ch}(G(n, \epsilon))$ is strictly contained in $H^{\text{even}}(G(n, \epsilon); \mathbb{Z})$ for each such group. These groups also afford p -group counterexamples to the conjecture of Atiyah described above.

Similar calculations may be made in the Brown-Peterson cohomology rings of these groups. These enable us to give a negative answer to a question of Landweber [7], who asked if Chern classes generate the Brown-Peterson cohomology of every p -group.

The examples.

The groups which we shall consider may be presented as

$$G(n, \epsilon) = \langle A, B, C \mid A^p = B^p = C^{p^{n-2}} = [B, C] = 1 \quad [A, C^{-1}] = B \quad [B, A] = C^{\epsilon p^{n-3}} \rangle,$$

where p is a prime not equal to 2 or 3, $n \geq 4$, and for fixed p and n there are two isomorphism classes of such groups, depending whether ϵ is either 1 or a quadratic non-residue modulo p . The group $G(n, \epsilon)$ has order p^n . In the sequel we shall refer to G instead of $G(n, \epsilon)$ unless the values of n and ϵ are important. The subgroup M generated by B and C is maximal (and hence normal) and is isomorphic to $C_p \oplus C_{p^{n-2}}$. We define one dimensional representations θ and ϕ of the group M by

$$\theta : B^j C^k \mapsto \exp(2\pi i j / p)$$

$$\phi : B^j C^k \mapsto \exp(2\pi i k / p^{n-2}).$$

The action of the quotient group G/M on the representation ring of M is that conjugation by A sends θ to $\theta \otimes \phi^{\otimes p^{n-3}}$ and sends ϕ to $\phi \otimes \theta^{\otimes \epsilon}$. Later we shall define elements of $H^2(M; \mathbb{Z})$ and $BP^2(BM)$ as Chern classes, and the action of G/M on these elements will be determined by its action on the representations θ and ϕ .

The group G has only 1- and p -dimensional irreducible representations because it has an abelian subgroup (M in fact) of index p . A one dimensional representation of G must

restrict trivially to $\langle B \rangle$, and a p -dimensional representation of G restricts to $\langle B \rangle$ as either p copies of the same representation of $\langle B \rangle$, or as the sum of one copy of each of the one-dimensional representations of $\langle B \rangle$. The examples $\text{Ind}_M^G(\theta)$ and $\text{Ind}_M^G(\phi)$ show that both these alternatives do occur.

Now define generators β, γ for $H^2(M; \mathbb{Z})$ by

$$\beta = c_1(\theta) \quad \gamma = c_1(\phi),$$

$$\text{so that} \quad H^{\text{even}}(M; \mathbb{Z}) \cong \mathbb{Z}[\beta, \gamma]/(p\beta, p^{n-2}\gamma),$$

and let β' be the restriction to $\langle B \rangle$ of β , so that

$$H^*(\langle B \rangle; \mathbb{Z}) \cong \mathbb{Z}[\beta']/(p\beta').$$

Lemma 1. *With notation as above, the image of $\text{Ch}(G)$ under restriction to $\langle B \rangle$ is the subring of $H^*(\langle B \rangle; \mathbb{Z})$ generated by β'^{p-1} and β'^p . For all $m \geq 0$,*

$$\beta'^{m+p-1} = -\text{Res}_{\langle B \rangle}^G \text{Cor}_M^G(\gamma^{p-1} \beta^m).$$

Proof. If ρ is a 1-dimensional representation of G then its Chern class restricts trivially to $\langle B \rangle$. If ρ is a p -dimensional representation of G , then either ρ restricts to $\langle B \rangle$ as p -copies of the same representation, in which case

$$\text{Res}(c.(\rho)) = (1 + i\beta')^p = 1 + i\beta'^p,$$

or as one copy of each representation, in which case

$$\text{Res}(c.(\rho)) = \prod_{i=0}^{p-1} (1 + i\beta') = 1 - \beta'^{p-1}.$$

By applying the double coset formula we see that

$$\begin{aligned} \text{Res}_{\langle B \rangle}^G \text{Cor}_M^G(\gamma^{p-1} \beta^m) &= \text{Res}_{\langle B \rangle}^M \left(\sum_{i=0}^{p-1} c_{A^i}^*(\gamma^{p-1} \beta^m) \right) \\ &= \text{Res}_{\langle B \rangle}^M \left(\sum_{i=0}^{p-1} (\gamma + i\epsilon\beta)^{p-1} (\beta + ip^{n-3}\gamma)^m \right) \\ &= -\beta'^{m+p-1} \end{aligned} \quad \blacksquare$$

Remarks. The image of $\text{Res}_{\langle B \rangle}^G$ is precisely the subring of $H^*(\langle B \rangle; \mathbb{Z})$ generated by $\beta'^{p-1}, \beta'^p, \beta'^{p+1}, \dots, \beta'^{2p-3}$. One way to show this is by considering the subgroup N of G generated by A and B . This subgroup is normal in G , and is the non-abelian group of order p^3 and exponent p . Using Lewis' calculation of $H^*(N; \mathbb{Z})$ [9], it may be shown that the image of $H^*(N; \mathbb{Z})^{G/N}$ under restriction to $\langle B \rangle$ does not contain β'^i for $i < p - 1$.

Corollary 2. $\text{Ch}(G)$ is strictly contained in $H^{\text{even}}(G; \mathbb{Z})$. Moreover, $\text{Ch}(G)$ does not map onto $H^*(G; \mathbb{Z})$ modulo its nilradical.

Proof. We know that β'^{p+1} is in $\text{Res}_{\langle B \rangle}^G(H^{\text{even}}(G; \mathbb{Z}))$, but not in $\text{Res}_{\langle B \rangle}^G(\text{Ch}(G))$. ■

Corollary 3. In the AHSS for G , write $B_\infty(G)$ for the universal boundaries, and $Z_\infty(G)$ for the universal cycles. Then $\text{Ch}(G) + B_\infty(G)$ is strictly contained in $Z_\infty(G)$.

Proof. The AHSS for $\langle B \rangle$ collapses, so $\text{Res}_{\langle B \rangle}^G(B_\infty(G))$ is trivial. Corestrictions of Chern classes are universal cycles, so $\beta'^{p+1} \in \text{Res}_{\langle B \rangle}^G(Z_\infty(G))$, but $\beta'^{p+1} \notin \text{Res}_{\langle B \rangle}^G(\text{Ch}(G) + B_\infty(G))$. ■

For any generalised cohomology theory \mathcal{H} and any group K , we may define $\text{Ch}_{\mathcal{H}}(K)$ to be the subring of $\mathcal{H}^*(BK)$ generated by $\rho^*(\mathcal{H}^*(BU))$ for all representations ρ of K in a unitary group U . We now give a result analogous to Corollary 2 for Brown-Peterson cohomology.

Lemma 4. $\text{Ch}_{BP}(G)$ is strictly contained in $BP^*(BG)$.

Proof. As in the integral cohomology case, define elements β and γ in $BP^2(M)$ by $\beta = c_1(\theta)$, $\gamma = c_1(\phi)$, and also define $\beta' = \text{Res}_{\langle B \rangle}^M(\beta)$, so that

$$BP^*(BM) \cong BP_*[[\beta, \gamma]]/([p]\beta, [p^{n-2}]\gamma), \quad BP^*(B\langle B \rangle) \cong BP_*[[\beta']]/([p]\beta'),$$

where $[r]x$ stands for the BP formal group sum of r copies of x . Let ' \equiv ' stand for congruence modulo the ideal of $BP^*(B\langle B \rangle)$ generated by p, v_1, v_2, \dots . As in Lemma 1, if ρ is a

p -dimensional representation of G , then either

$$\begin{aligned} \text{Res}_{\langle B \rangle}^G(c.(\rho)) &= (1 + [i]\beta')^p \equiv 1 + i\beta'^p \\ \text{or } \text{Res}_{\langle B \rangle}^G(c.(\rho)) &= \prod_{i=0}^{p-1} (1 + [i]\beta') \equiv 1 - \beta'^{p-1}. \end{aligned}$$

Also, we have that

$$\begin{aligned} \text{Res}_{\langle B \rangle}^G \text{Cor}_M^G(\gamma^{p-1}\beta^2) &= \text{Res}_{\langle B \rangle}^M \left(\sum_{i=0}^{p-1} (\gamma +_{BP} [i\epsilon]\beta)^{p-1} (\beta +_{BP} [ip^{n-3}]\gamma)^2 \right) \\ &\equiv -\beta'^{p+1}. \end{aligned} \quad \blacksquare$$

Our original proofs of these results involved calculation with $BP^*(BN)$, which has been determined by Tezuka-Yagita [10], and with the integral cohomology of the non-abelian maximal subgroups of G , determined by Leary [8]. Using these methods we obtain more information concerning $BP^*(BG)$ and $H^*(G; \mathbb{Z})$, which we intend to publish later.

We thank Dr. C. B. Thomas for suggesting investigations of the Chern subring and for his encouragement. The second author also thanks M. Tezuka and M. Kaneda for many suggestions. The work of the first author has been supported by the SERC.

REFERENCES

- [1] K. ALZUBAIDY. Metacyclic p -groups and Chern classes. *Illinois J. Math.* **26** (1982), 423–431.
- [2] K. ALZUBAIDY. Rank 2 p -groups, $p > 3$, and Chern classes. *Pacific J. Math.* **103** (1982), 259–267.
- [3] M. F. ATIYAH. Characters and the cohomology of finite groups. *Publ. Math. IHES* **9** (1961), 23–64.
- [4] N. BLACKBURN. Generalisations of certain elementary theorems on p -groups. *Proc. London Math. Soc.* **11** (1961), 1–22.
- [5] K. S. BROWN. *Cohomology of groups*. Springer Verlag, (1982).

- [6] J. HUEBSCHMANN. Chern classes for metacyclic groups. preprint (1988).
- [7] P. LANDWEBER. Elements of infinite filtration in complex cobordism. *Math. Scand.* **30** (1972), 223–6.
- [8] I. J. LEARY. The integral cohomology rings of some p -groups. Submitted.
- [9] G. LEWIS. The integral cohomology rings of groups of order p^3 . *Trans. Amer. Math. Soc.* **132** (1968), 501–529.
- [10] M. TEZUKA and N. YAGITA. Cohomology of finite groups and Brown-Peterson cohomology. *Springer LNM* **1370** (1989), 396–408.
- [11] M. TEZUKA and N. YAGITA. Cohomology of finite groups and Brown-Peterson cohomology II. *Springer LNM* **1418** (1990), 57–69.
- [12] C. B. THOMAS. Chern classes and metacyclic p -groups. *Mathematika* **18** (1971), 196–200.
- [13] C. B. THOMAS. Riemann-Roch formulae for group representations. *Mathematika* **20** (1973), 253–262.
- [14] C. B. THOMAS. Filtrations on the representation ring of a finite group. *Cont. Math.* **19** (1983), 397–405.
- [15] C. B. THOMAS. *Characteristic classes and the cohomology of finite groups*. Cambridge University Press, (1986).
- [16] E. A. WEISS. Bonner Mathematische Schriften. 1969